

A helium-hydrogenic molecular atmosphere of neutron star 1E1207.4-5209

Alexander V. Turbiner*

*Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México,
Apartado Postal 70-543, 04510 México, D.F., México.*

Abstract

A model of a mixed helium-hydrogenic atmosphere of the isolated neutron star 1E1207.4-5209 is proposed. It is based on the assumption that the main components in the atmosphere are the exotic molecular ions He_2^{3+} and H_3^{2+} with the presence of He^+ , $(HeH)^{2+}$, H_2^+ under a surface magnetic field $\approx 4.4 \times 10^{13}$ G. In addition to two absorption features observed by *Chandra* observatory (Sanwal et al, 2002) the model predicts one more narrow absorption feature at ≈ 400 eV.

PACS numbers: 31.15.Pf, 31.10.+z, 32.60.+i, 97.10.Ld

arXiv:astro-ph/0506677v1 27 Jun 2005

*On leave of absence from the Institute for Theoretical and Experimental Physics, Moscow 117259, Russia;

Electronic address: turbiner@nucleares.unam.mx

Recently, it was discovered by Chandra *X-ray* observatory the absorption features at 0.7 KeV and 1.4 KeV in the *X-ray* spectrum of the isolated neutron star 1E1207.4-5209 [1] (see Fig.1). Later this discovery was confirmed by XMM-Newton *X-ray* observatory [2]. These features had widths 0.1 and 0.13 KeV, respectively. For many people it was quite impressive property that the second absorption energy is about twice larger than the first one. Different proposals about content of the atmosphere are presented. All of them are agreed on partially ionized atmosphere content. Almost all of them are related to atomic ions (for a review, see [1, 3]). Among of these models it was proposed a model of the hydrogenic atmosphere of 1E1207.4-5209 with main abundance of the exotic ion H_3^{2+} [4]. In particular, this model explains the observed absorption features as a consequence of the photodissociation $H_3^{2+} \rightarrow H + 2p$ and the H_3^{2+} photoionization, while the bound-bound transitions are assumed to be non-important. However, this model requires very strong magnetic field on the surface, $(2 - 6) \times 10^{14}$ G.

Above-mentioned discovery pushed us for searching new exotic chemical compounds in a strong magnetic field in addition to H_3^{2+} [5]. Very recently, we had discovered that in the domain $4.4 \times 10^{13} \gtrsim B \gtrsim 10^{11}$ G two more exotic systems can occur: the molecular ions $(\text{HeH})^{2+}$ and He_2^{3+} [6]. These exotic systems do not exist without magnetic field. Among these two the one-electron molecular ion $\text{He}e_2^{3+}$ [6] plays exceptionally important role at large magnetic fields: at $B \gtrsim 3 \times 10^{12}$ G it becomes the Coulomb system with lowest total energy among one-electron systems made from protons and/or α -particles. It was also quite surprising to find out that at the same domain of magnetic fields another exotic ion H_3^{2+} becomes the Coulomb system with lowest total energy among one-electron systems made from protons. It is worth emphasizing that in very large domain of magnetic fields, $2 \times 10^{14} \gtrsim B \gtrsim 2 \times 10^{12}$ G the ratio of the binding (ionization) energies of these two systems remains almost constant,

$$\frac{E_T^{H_3^{2+}}}{E_T^{\text{He}e_2^{3+}}} \approx 0.5 .$$

In this Note we propose a mixed helium-hydrogenic model of atmosphere of the 1E1207.4-5209. This model provides an explanation of both absorption features and requires a surface magnetic field $\approx 4 \times 10^{13}$ G, which is not in so striking contradiction with that estimated using the period derivative, $(2 - 4) \times 10^{12}$ G (see [1]).

Let us make the simplest assumption that the atmosphere consists of α - particles, pro-

tons and electrons mostly in a form of the H , H_2^+ , $H_3^{(2+)}$, $H_4^{(3+)}$, $(\text{HeH})^{2+}$, He^+ and He_2^{3+} systems [9]. The first important observation is the fact that the only neutral system H is always characterized the highest total energy among above-mentioned systems for any chosen magnetic field strength [5, 6]. Hence, in general, it can be neglected. It is evident that the charged systems H_2^+ , $H_3^{(2+)}$, $H_4^{(3+)}$, $(\text{HeH})^{2+}$, He^+ and He_2^{3+} can move mostly in the longitudinal (along the magnetic line) direction, and their transverse motion is limited to a domain defined by a Larmor radius. They are divided into three types: ionization (bound-free transitions), dissociation and excitation (bound-bound transitions). A study of all above-mentioned systems were done using the variational technique with physically relevant trial functions (see e.g. [5, 6]. In Table I the binding energies for magnetic field 4.4×10^{13} G are given.

It can be immediately seen that the binding (ionization) energies of the most bound one-electron system H_3^{2+} as well as H_2^+ belong to the domain of the first absorption feature at 0.7 KeV. While the binding (ionization) energies of the $(\text{HeH})^{2+}$, He^+ and He_2^{3+} systems correspond to the second absorption feature at 1.4 KeV. In Table II we present the disso-

TABLE I: *Binding energies in Rydbergs (Ry) and in electron-volts (eV) for one-electron Coulomb systems for magnetic field 4.4×10^{13} G. Energies in eV are rounded to the nearest integer number ending in 0 or 5.*

H -atom	He^+	He_2^{3+}	$(\text{HeH})^{2+}$	H_2^+	H_3^{2+}	
33.8	93.5	105.1	92.6	54.5	55.2	Ry
460	1270	1430	1260	740	750	eV

ciation energies. The range of sensitivity of the *Chandra/ACIS* detector does not allow to see the domain where the photodissociation process $\text{He}_2^{3+} \rightarrow \text{He}^+ + \alpha$ can contribute. In principle, two other photodissociation processes $H_3^{2+} \rightarrow H + 2p$ and $H_2^+ \rightarrow H + p$ can be seen although they are close to the end of the range of sensitivity of the *Chandra/ACIS* detector.

In Table III we present the energies of the first excitation of H_2^+ and H_3^{2+} , correspondingly. These processes contribute to the domain which is close to the end of the range of sensitivity of the *Chandra/ACIS* detector and unlikely to be detected. The first excitation of He_2^{3+} can be seen. This line can be quite narrow. Hence, the domain around 0.4 KeV deserves to be

TABLE II: *Dissociation energies in Rydbergs (Ry) and electron-volts (eV) for some one-electron systems for magnetic field 4.4×10^{13} G. Energies in eV are rounded to the nearest integer number ending in 0 or 5.*

$He_2^{3+} \rightarrow He^+ + \alpha$	$H_2^+ \rightarrow H + p$	$H_3^{2+} \rightarrow H + 2p$	
11.6	20.7	21.4	Ry
160	280	290	eV

TABLE III: *Excitation energies in Rydbergs (Ry) and electron-volts (eV) for some one-electron systems for magnetic field 4.4×10^{13} G. Energies in eV are rounded to the nearest integer number ending in 0 or 5.*

$He_2^{3+}(1\sigma_g \rightarrow 1\pi_u)$	$H_2^+(1\sigma_g \rightarrow 1\pi_u)$	$H_3^{2+}(1\sigma_g \rightarrow 1\pi_u)$	
29.0	13.4	15.0	Ry
395	180	205	eV

observed carefully. We assume that the photoexcitation cross-sections are smaller than the photoionization ones following the results obtained for the hydrogen atom [7].

We can formulate our model of the content of the neutron star atmosphere. It corresponds to the surface magnetic field $\approx 4.4 \times 10^{14}$ G. The atmosphere is partially ionized and consists of H_2^+ , $H_3^{(2+)}$, He^+ , $(HeH)^{2+}$, He_2^{3+} ions. Photoionization of H_2^+ , H_3^{2+} explains the first absorption feature in Fig.1 at 0.7 KeV while photoionization He^+ , $(HeH)^{2+}$, He_2^{3+} explains the second absorption feature at 1.4 KeV [10]. We predict a narrow absorption feature emerging from electronic excitation of $He_2^{3+}(1\sigma_g \rightarrow 1\pi_u)$ around 400 KeV. It can allow to identify the present model. Red-shift effects are not taken into account, but they can be included straightforward (see e.g. [1]) and we guess can increase the magnetic field values for $\sim 20 - 50\%$. We think that the presence of the gravitational sedimentation of hydrogen and helium will not change our results.

The model of the atmosphere with main abundance of the He^+ atomic ions proposed in [1] with the surface magnetic field 1.5×10^{14} G is lacking the explanation why a possible presence of the molecular ion He_2^{3+} is neglected. For such a magnetic field the molecular ion He_2^{3+} is much more bound than the atomic ion He^+ . It can be calculated [11] that total




FIG. 1: *Chandra*/ACIS spectrum as presented in [3] (Fig.#1 from [3] by the author's permission). The solid line is a blackbody model of [3] which is corrected by the detector sensitivity to illustrate two absorption features.

energy of He_2^{3+} at $B = 1.5 \times 10^{14}$ G is lower than the total energy of He^+ to ≈ 26.4 Ry. Even under NS surface temperature 1.4-1.9 MK (see, e.g. [1]) it is not clear why the presence of He_2^{3+} should be discarded. Perhaps, it is worth mentioning that for the magnetic field 1.5×10^{14} G another exotic molecular ion $(HeH)^{2+}$ is also more bound than He^+ :

$$E_T^{He^+} - E_T^{(HeH)^{2+}} \approx 4.4 \text{ Ry} .$$

The author is grateful to J.C. Lopez Vieyra. K. Mori, G.G. Pavlov and A.Y. Potekhin for valuable discussions. The work is supported in part by CONACyT grant 47899-E.

-
- [1] D. Sanwal et al., ‘Discovery of absorption features in the X-ray spectrum of an isolated neutron star’,
ApJL, **574**, L61 (2002)
(astro-ph/0206195)
 - [2] G.F. Bignami, P.A. Caraveo, A. De Luca and S. Mereghetti, *Nature* **423**, 725-727 (2003)
 - [3] C.J. Hailey and K. Mori, ‘Evidence for a mid-atomic-number atmosphere in the neutron star 1E1207.4-5209’,
ApJL, **578**, L133 (2002)
(astro-ph/0206195)

K. Mori and C.J. Hailey, ‘X-ray spectroscopy of the isolated neutron star 1E1207.4-5209: atmospheric composition and equation of state’
(astro-ph/0301161)
 - [4] A.V. Turbiner and J.-C. Lopez Vieyra, *Mod.Phys.Lett.* **A19**, 1919-1923 (2004)
(astro-ph/0404290)
 - [5] A. Turbiner, J.-C. Lopez and U. Solis H., Predicted existence of H_3^{++} molecular ions in strong magnetic fields, *Pisma v ZhETF* **69**, 800-805 (1999); *JETP Lett.* **69**, 844-850 (1999) (English Translation), (astro-ph/980929)
 - [6] A.V. Turbiner and J.-C. Lopez Vieyra, Preprint ICN-UNAM 04-15, pp.7, December 2004
(astro-th/0412399)
 - [7] A.Y. Potekhin and G. Chabrier, *ApJ* **600**, 317 (2004)
(astro-ph/0309310)
 - [8] D. Lai, E. Salpeter and S.L. Shapiro, *Phys. Rev.* **A45**, 4832 (1992);
D. Lai and E. Salpeter, *Phys. Rev.* **A53**, 152 (1996);
D. Lai, ‘Matter in Strong Magnetic Fields’, *Rev. Mod. Phys.* **73**, 629 (2001)
(astro-ph/0009333)
 - [9] In principle, deuterons or tritons can be instead of protons.
 - [10] The cross-section of photo-ionization depends on the energy of ionization. We unaware of any reliable calculations of bound-bound and bound-free transitions, even for the simplest molecular system H_2^+ , we assume, following a detailed study of the hydrogen atom [7], that

(i) photo-ionization cross-section has a maximum near the ionization threshold, and (ii) bound-bound transition amplitudes are small compared to the bound-free ones. We also neglect any difference between ionization threshold (binding energy) and the maximum of the energy distribution, assuming that this difference is small.

- [11] In non-relativistic framework, neglecting the center-of-mass effects and relativistic corrections, which we assume following the Salpeter et al (see [8]) do not exceed 10% in the binding energy.